



Potential insecticidal toxins extracted from some isolates of *Akanthomyces*

T. Soltani ¹, F. Yarahmadi ^{2*}, A. Rajabpour ², M.H. Ghodoum Parizipour ³

1. M.Sc. Graduate of Agricultural Entomology, Department of Plant Protection, Faculty of Agriculture, Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani, Iran
2. ***Corresponding Author:** Professor, Department of Plant Protection, Faculty of Agriculture, Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani, Iran (fa_yarahmadi@yahoo.com)
3. Assistant Professor, Department of Plant Protection, Faculty of Agriculture, Agricultural Sciences and Natural Resources University of Khuzestan, Mollasani, Iran

Received: 1 September 2023

Accepted: 25 November 2023

Abstract

Entomopathogenic fungi (EPF) are key elements for pest management in sustainable agriculture. Many EPFs can produce secondary metabolites with toxic effects, playing an essential role in their pathogenicity. In this study, the metabolites of some EPFs, including three isolates of *Akanthomyces lecanii* species (PAL6, PAL7, and PAL8) and one isolate of *A. muscarius* species (AGM5), were biochemically analyzed. Extracellular metabolites were isolated from the fungal cells of the two-week-old culture of each isolate, and the chemical constituents were identified by high-performance liquid chromatography (HPLC). During the retention interval of 10-13 minutes, four different peaks were observed within metabolites of the isolates PAL6, PAL7, and PAL8. In the case of the AGM5 isolate; however, one peak was observed at 11.53 retention time. Comparative analysis showed the presence of insecticidal toxic cyclic peptides such as bassianolide within the fungal metabolites. Moreover, one peak with a new identity was detected within the metabolites from the fungal isolates of PAL6, PAL7, and PAL8 in the retention interval of 5 minutes from the isolate AGM5 at 11.53 retention times. The results demonstrated that producing toxic compounds such as bassianolide by the EPF probably contributed to their insecticidal effects.

Keywords: *Mycotoxins; cyclic peptides; bioinsecticide; entomopathogenic fungi*

Associate editor: M. Yazdani (Ph.D.)

Citation: Soltani, T., Yarahmadi, F., Rajabpour, A. & Ghodoum Parizipour, M.H. (2023). Potential Insecticidal Toxins Extracted from Some Isolates of *Akanthomyces*. *Plant Protection (Scientific Journal of Agriculture)*, 46(3), 17-23. <https://doi.org/10.22055/ppr.2023.44689.1706>.

Introduction

Many entomopathogenic fungi (EPFs), as the most important microbial control agents of insect pests, systematically belong to the phylum Ascomycota, order Hypocreales, and family Cordycipitaceae (Litwin et al., 2020). The genus *Akanthomyces* is an important EPF from the family Cordycipitaceae (Ascomycota, Hypocreales), infecting spiders and insects in various ecosystems (Aini et al., 2020). The genus that was first established by Lebert (1858) can parasitize many insect orders, including Hemiptera, Coleoptera, Lepidoptera and Orthoptera (Mongkolsamrit et al., 2018). Among the genera, *A. lecanii* Zimmerman and *A. muscarius* Petch are two important and widespread EPFs of hemipteran pests (Broumandnia et al., 2021). Three Iranian-originated isolates of *A. lecanii* (PAL6, PAL7, and PAL8) and one isolate of *A. muscarius* (AGM5) were previously isolated from hemipteran citrus pests, *Pulvinaria aurantii* Cock. (Hemiptera: Coccidae) and *Aphis gossypii* Glover (Hemiptera: Aphididae). The appropriate pathogenicity of the isolates against *Bemisia tabaci* Gen. (Hem., Aleyrodidae) (Broumandnia et al., 2021), *Tribolium castaneum* Herbst (Col., Tenebrionidae) (Broumandnia & Rajabpour, 2020) and *Aphis fabae* Scopoli (Hemiptera: Aphididae) (Soltani et al., 2022) have been previously reported.

Similar to other EPFs, *Akanthomyces* fungi can produce some secondary metabolites that are considered mycotoxins. Some metabolites have considerable insecticidal effects and lead to higher and faster insect host mortality in comparison with nontoxicogenic fungi (Wang et al., 2018). Furthermore, the antifeedant properties of some active metabolites of EPF have been documented (Berestetskiy & Hu, 2021). In some cases, metabolites induce host plant systematic resistance to sap-feeding plants, e.g., whiteflies, by triggering related genes (Abdulle et al., 2021). In total, secondary metabolites play important roles in host plant resistance against insect pests. Some

bioinsecticides have been developed based on mycotoxin constituents derived from EPFs, e.g., beauvericin from *Beauveria bassiana* Balsamo and cytochalasin C from *Metarhizium anisopliae* Metschn. and *Tolypocladium inflatum* Gams (Bogus et al., 2021). Due to many problems and limitations associated with chemical insecticides, bioinsecticides can play an important role in the eco-friendly control of insect pests for developing integrated pest management (IPM) programs. Determinations of mycotoxin constituents are the basic information before bioinsecticide production. The chemical compounds of some mycotoxins derived from *Acanthomyces* fungi were previously studied. For instance, constituents of bioactive metabolites extracted from mycelia of *A. evansii* comprised phenolic compounds (terphenylin, deoxy terphenylin, terprenin 2, terprenin epoxide), bipeptide (cyclo-tyrosylprolyl), and simple aromatic compounds (acetyl hydroxybenzamide, 4-hydroxybenzaldehyde) (Effendi, 2012). It has been reported that many of the bioactive metabolites derived from *Akanthomyces* act as cuticle-degrading proteases to facilitate penetration of EPF mycelia (Keppanan et al., 2019). This study aimed to detect the insecticidal bioactive compounds derived from pathogenic isolates of *Akanthomyces*.

Materials and methods

Fungal culture and maintenance

Different isolates of pathogenic fungi from *P. aurantii* and *A. gossypii* were isolated in citrus orchards of northern Iran, 36°54'24.2"N 50°39'26.7"E, and their pathogenicity was confirmed against *T. castaneum* (Broumandnia and Rajabpour, 2020), *B. tabaci* (Broumandnia et al., 2021) and *A. fabae* (Soltani et al., 2022). Morphological and molecular investigations have indicated that EPFs belong to three fungal isolates of *A. lecanii* species (PAL6, PAL7 and PAL8) and one fungal isolate of *A. muscarius* species (AGM5) (Broumandnia and Rajabpour, 2020; Broumandnia et al., 2021). The isolates were cultured on potato dextrose agar (PDA)

medium in a germinator with a temperature of $23\pm 1^\circ\text{C}$, relative humidity of 43%, and 16:8 light: dark hours.

Preparation of crude secondary metabolites

Mycelia from the two-week-old culture of the fungi were harvested and isolated using centrifugation at 1000 g. The bioactive compounds were isolated from the fungi after an incubation period of two weeks. Extracellular metabolites were isolated from the fungal cells using methanol:chloroform (1:2, v/v). Then, the mixture was centrifuged at 1000 g to remove fatty acids. The organic phase was separated and concentrated in a vacuum rotary evaporator. Relatively purified insecticidal toxic compounds were recovered and analyzed.

High-performance liquid chromatography (HPLC)

The crude extract of the fungal isolates was dissolved in methanol-acetonitrile solution. The mobile phase was run in a K-60 silica gel column (230–400 mesh, 325 240 mm, Darmstadt, Germany) of chromatography machine (KNAUER, Germany) and started step by step in 50-mm methylene. First, methylene:methanol-dichloride (95:5) was used, and the slope of this solvent was used to achieve the desired polarities. The prepared sample was diluted with methanol and analyzed using a reversed-phase HPLC column, and the peaks were detected by absorbing ultraviolet light at a wavelength of 240 nm. A typical slope of the thigh was used as follows: 0 minutes (0% acetonitrile), 30 minutes (40% acetonitrile), 40 minutes (50% acetonitrile), and 60 minutes (50% acetonitrile).

Results and discussion

Fungal metabolites were subjected to HPLC analysis by a reversed-phase C18 column, and the compounds were identified. The analysis report showed the presence of the insecticide's toxic cyclic peptide, e.g., bassianolide. Among the retention times of 10-13 minutes, four different peaks were observed for isolates PAL6, PAL7 and PAL8 (Figure 1).

In this way, these peaks in the sample obtained from the PAL6 isolate at retention times of 10.85, 11.97 and 12.35, the samples belonging to PAL7 at retention times of 11.12, 12.27 and 12.65, and samples related to PAL8 appeared at retention times of 10.53, 11.53 and 11.87. Three peaks with a new identity were detected within the metabolites from the fungal isolates of PAL6, PAL7 and PAL8 at a retention time of 5 minutes. In addition, other peaks were also observed in the chromatograms of these samples, which were not considered due to the time interval from the target intervals. In the case of the GMA5 isolate, however, one peak was observed at 11.53 retention time. In addition, a peak with high intensity also appeared at a retention time of 5.48. Comparison of the peaks with those of insecticidal compounds (Ravindran et al., 2018) revealed the presence of toxic compounds within the metabolites of tested isolates. However, the low intensity of the peaks reflects the relatively low concentration of these compounds. Furthermore, the peaks observed at 5 minutes, especially for the GMA5 isolate, can indicate the presence of a new compound in the fungal metabolites of these isolates, which requires additional tests in this regard.

EPF secrete various nonvital compounds that are key elements for infection to infect their hosts (Vey et al., 2001). Destruxins (A, B, C, and E) are a major group of compounds identified from *M. anisopliae* (Ravindran et al., 2016). *Akanthomyces* spp. are famous EPFs used for pest biocontrol, and their insecticidal metabolites have been applied against various insects (Butt et al., 1994; Isaka et al., 2005; Molnar et al., 2010). Among the genera, *A. lecanii* is an EPF that has been well studied as a microbial biocontrol agent, and the compounds derived from the fungus have appropriate toxic effects and have been applied against various arthropodan pests (Butt et al., 1994; Isaka et al., 2005; Molnar et al., 2010). Many compounds with insecticidal properties produced by the pathogen are cyclic peptides (Ravindran et al., 2018). Bassianolide is a toxic compound that has been identified within

A. lecanii metabolites (Ravindran et al., 2018). Bassianolide disrupts insect smooth muscle contraction by inhibiting the acetylcholine mechanism. Furthermore, the compound has cytotoxic, moderate antiplasmodial and antimycobacterial activities (Jirakkakul et al., 2008). In vitro studies indicated that bassianolide significantly induces the virulence of EPF, e.g., *B. bassiana*, to the larvae of *Helicoverpa zea* Boddie (Lepidoptera; Noctuidae) and *Bombyx mori* L. (Lepidoptera: Bombycidae). Therefore, the compound serves as a bona fide virulence factor of the EPF, significantly contributing to the commercial microbiological insecticide preparation containing the fungus conidia (Singh et al., 2015). Similarly, bassianolide was detected within metabolites of Iranian isolates of

A. lecanii and *A. muscarius*. EPF usage against arthropodian hosts makes them more vulnerable to pathogen reinvasion. However, knowledge about the mechanisms of insect host defense and its specific bioresponse when it is exposed to metabolites is needed (Wu et al., 2016). The crude extract from the Iranian isolates of *A. lecanii* and *A. muscarius* showed the presence of two major and four minor peaks (Figures 1), which are characteristic of bassianolide. Previous studies have also characterized the presence of cyclic peptides such as beauvericin, bassianolide and DXTs from EPFs such as *B. bassiana* and *M. anisopliae* using HPLC analysis (Kershaw et al., 1999; Liu et al., 2009; Namara et al., 2017; Ravindran et al., 2016; Sowjanya Sree et al., 2008).




Figure 1. Chromatograms of metabolites of *Akanthomyces* isolates, including PAL7 (a), PAL6 (b) GMA5 (c) and PAL8 (d), using HPLC.

Our results demonstrated that bassianolide probably contributes to the insecticidal effects of *A. lecanii* *A. muscarius* against pests such as *T. castaneum* (Broumandnia & Rajabpour, 2020), *B. tabaci* (Broumandnia et al., 2021) and *A. fabae* (Soltani et al., 2022) and can be a promising source of toxic metabolites against other pests.

Conclusion

Both EPFs, *A. lecanii* and *A. muscarius*, are well-known EPFs that have been applied against several insect pests. They can produce secondary metabolites with toxic effects by which they

fulfill their pathogenicity process. HPLC analysis of the metabolites of the EPF showed the presence of insecticidal toxic cyclic peptides such as bassianolide within the fungal metabolites. Moreover, one peak with a new identity was detected within the metabolites from the EPF. These toxic compounds are probably responsible for the insecticidal effects of the EPF.

Acknowledgments

The authors thank Dr. S. Aghajanzadeh and Citrus and Subtropical Fruits Research Center of Iran for their technical assistance.

References

- Abdulle, Y. A., Nazir, T., Sayed, S., Mahmoud, S. F., Majeed, M. Z., Aslam, H. M. U., & Qiu, D. (2021). Sub-Lethal Effects of *Lecanicillium lecanii* (Zimmermann)-Derived Partially Purified Protein and Its Potential Implication in Cotton (*Gossypium hirsutum* L.) Defense against *Bemisia tabaci* Gennadius (Aleyrodidae: Hemiptera). *Agriculture*, 11(8), 778. DOI: <https://doi.org/10.3390/agriculture11080778>
- Aini, A. N., Mongkolsamrit, S., Wijanarka, W., Thanakitpipattana, D., Luangsa-Ard, J. J., & Budiharjo, A. (2020). Diversity of *Akanthomyces* on moths (Lepidoptera) in Thailand. *MycKeys*, 71, 1. DOI: [10.3897/mycokeys.71.55126](https://doi.org/10.3897/mycokeys.71.55126).
- Azadi, F., Rajabpour, A., Lotfi Jalal Abadi, A., & Mahjoub, M. (2018). Resistance of tomato cultivars to *Tuta absoluta* (Lepidoptera: Gelechiidae) under field condition. *Journal of Crop Protection*, 7(1), 87-92.
- Berestetskiy, A. and Hu, Q. (2021). The chemical ecology approach to reveal fungal metabolites for arthropod pest management. *Microorganisms*, 9(7), 1379. DOI: <https://doi.org/10.3390/microorganisms9071379>
- Boguś, M. I., Wrońska, A. K., Kaczmarek, A., & Boguś-Sobocińska, M. (2021). In vitro screening of 65 mycotoxins for insecticidal potential. *Plos one*, 16(3), e0248772.
- Broumandnia, F., & Rajabpour, A. (2020). Efficacies of some isolates of *Lecanicillium lecanii* to control *Tribolium castaneum* (Col., Tenebrionidae). *Journal of Plant Diseases and Protection*, 127(5): 625-631. DOI: <https://doi.org/10.1007/s41348-020-00324-y>.
- Broumandnia, F., Rajabpour, A., Parizipour, M. H. G., & Yarahmadi, F. (2021). Morphological and molecular identification of four isolates of the entomopathogenic fungal genus *Akanthomyces* and their effects against *Bemisia tabaci* on cucumber. *Bulletin of Entomological Research*, 111(5), 628-636. DOI: <https://doi.org/10.1017/S0007485321000298>.
- Butt, T. M., Ibrahim, L., Ball, B. V., & Clark, S. J. (1994). Pathogenicity of the entomogenous fungi *Metarhizium anisopliae* and *Beauveria bassiana* against crucifer pests and the honey bee. *Biocontrol Science and Technology*, 4, 207-214. DOI: <https://doi.org/10.1080/09583159409355328>.

- Effendi, H. (2012). Phenolic Compounds of Sponge-associated Fungi (*Lecanicillium evansii*). *Microbiology Indonesia*, 6(3): 1-1.
- Isaka, M., Kittakoop, P., Kirtikara, K., Hywel-Jones, N.L., & Thebtaranonth, Y. (2005). Bioactive substances from insect pathogenic fungi. *Acc. Chemical Research*, 38, 813–823. DOI: <https://doi.org/10.1021/ar040247r>.
- Jirakkakul, J., Punya, J., Pongpattanakitsote, S., Paungmoung, P., Vorapreeda, N., Tachaleat, A., & Cheevadhanarak, S. (2008). Identification of the nonribosomal peptide synthetase gene responsible for bassianolide synthesis in wood-decaying fungus *Xylaria* sp. BCC1067. *Microbiology*, 154, 995–1006. DOI: <https://doi.org/10.1099/mic.0.2007/013995-0>.
- Keppanan, R., Sivaperumal, S., Hussain, M., Bamisile, B. S., Aguila, L. C. R., Qasim, M., & Krutmuang, P. 2019. Molecular characterization of pathogenesis involving the GAS 1 gene from Entomopathogenic fungus *Lecanicillium lecanii* and its virulence against the insect host *Diaphorina citri*. *Pesticide Biochemistry and Physiology*, 157, 99-107. DOI: <https://doi.org/10.1016/j.pestbp.2019.03.012>.
- Kershaw, M.J., Moorhouse, E.R., Bateman, R., Reynolds, S.E., & Charnley, A.K. (1999). The role of destruxins in the pathogenicity of *Metarhizium anisopliae* for three species of insect. *Journal of Invertebrate Pathology*, 74, 213–223. DOI: <https://doi.org/10.1006/jipa.1999.4884>.
- Lebert H. (1858). Über einige neue oder unvollkommen gekannte Krankheiten der Insekten, welche durch Entwicklung niederer Pflanzen im lebenden Körper entstehen. *Zeitschrift für wissenschaftliche Zoologie*, 9, 439–453.
- Litwin, A., Nowak, M., & Różalska, S. (2020). Entomopathogenic fungi: unconventional applications. *Reviews in Environmental Science and Biotechnology*, 19, 23–42. DOI: <https://doi.org/10.1007/s11157-020-09525-1>.
- Liu, W.M., Xie, Y.P., Xue, J.L., Gao, Y., Zhang, Y.F., & Jun, W. (2009). Histopathological changes of Ceroplastes japonicus infected by *Lecanicillium lecanii*. *Journal of Invertebrate Pathology*, 101: 96–105. DOI: <https://doi.org/10.1016/j.jip.2009.03.002>.
- Mohammadi, S., Seraj, A. A., & Rajabpour, A. (2015). Evaluation of six cucumber greenhouse cultivars for resistance to *Tetranychus turkestanii* (Acari: Tetranychidae). *Journal of Crop Protection*, 4(4), 545-556.
- Molnar, I., Gibson, D.M., & Krasnoff, S.B. (2010). Secondary metabolites from entomopathogenic Hypocrealean fungi. *Natural Product Report*, 27, 1241–1275. DOI: <https://doi.org/10.1039/C001459C>.
- Mongkolsamrit, S., Noisripoom, W., Thanakitpipattana, D., Wuthikun, T., Spatafora, J.W., & Luangsa-ard, J.J. (2018). Disentangling cryptic species with isaria-like morphs in Cordycipitaceae. *Mycologia*, 110(1), 230–257. DOI: <https://doi.org/10.1080/00275514.2018.1446651>.
- Namara, L.M., Carolan, J.C., Griffin, C.T., Fitzpatrick, D., & Kavanagh, K. (2017). The effect of entomopathogenic fungal culture filtrate on the immune response of the greater wax moth, *Galleria mellonella*. *Journal of Insect Physiology*, 100, 82–92. DOI: <https://doi.org/10.1016/j.jinsphys.2017.05.009>.

Ravindran, K., Akutse, K.S., Sivaramakrishnan, S., & Wang, L. (2016). Determination and characterization of destruxin production in *Metarhizium anisopliae* Tk6 and formulations for *Aedes aegypti* mosquitoes control at the field level. *Toxicon*, 120, 89–96. DOI: <https://doi.org/10.1016/j.jinsphys.2017.05.009>.

Ravindran, K., Sivaramakrishnan, S., Hussain, M., Dash, C. K., Bamisile, B. S., Qasim, M., & Liande, W. 2018. Investigation and molecular docking studies of Bassianolide from *Lecanicillium lecanii* against *Plutella xylostella* (Lepidoptera: Plutellidae). *Comparative Biochemistry and Physiology Part C: Toxicology and Pharmacology*, 206, 65-72. DOI: <https://doi.org/10.1016/j.cbpc.2018.03.004>.

Ren, S.X., Ali, S., Huang, Z., & Wu, J.H. (2010). *Lecanicillium muscarium* as microbial insecticide against whitefly and its interaction with other natural enemies. *Microbiology and Microbial Biotechnology*, 27, 339-348.

Singh, H. B., Keswani, C., Ray, S., Yadav, S. K., Singh, S. P., Singh, S., & Sarma, B. K. (2015). *Beauveria bassiana*: biocontrol beyond lepidopteran pests. In *Biocontrol of Lepidopteran Pests* (pp. 219-235). Springer, Cham.

Soltani, T., Yarahmadi, F., Rajabpour, A., & Ghoddom Parizi Pour, M. H. (2022). Pathogenicity of Iranian isolates of *Akanthomyces lecanii* and *A. muscarius* on the black bean aphid (*Aphis fabae* Scopoli). *Plant Protection (Scientific Journal of Agriculture)*, 45(1), 19-28. DOI: 10.22055/PPR.2021.17246.

Sowjanya Sree, K., Padmaja, V., & Murthy, Y.L.N. (2008). Insecticidal activity of destruxin, a mycotoxin from *Metarhizium anisopliae* (Hypocreales), against *Spodoptera litura* (Lepidoptera: Noctuidae) larval stages. *Pest Management Science*, 64, 119–125. DOI: <https://doi.org/10.1002/ps.1480>.

Vey, A., Hoagland, R.E., & Butt, T.M. (2001). Toxic metabolites of fungal biocontrol agents. In: Butt, T.M., Jackson, C.W., Magan, N. (Eds.), *Fungi as Biocontrol Agents: Progress, Problems and Potential*. CABI Publishing, Wallingford, UK, pp. 311–346.

Wang, X., Gong, X., Li, P., Lai, D., & Zhou, L. (2018). Structural diversity and biological activities of cyclic Depsipeptides from fungi. *Molecules*, 23,169. DOI: <https://doi.org/10.3390/molecules23010169>.

Wu, S., Xie, H., Li, M., Xu, X., & Lei, Z. (2016). Highly virulent *Beauveria bassiana* strains against the two-spotted spider mite, *Tetranychus urticae* show no pathogenicity against five phytoseiid mite species. *Experimental and Applied Acarology*, 70, 421–435. DOI: <https://doi.org/10.3390/molecules23010169>.



© 2023 by the authors. Licensee SCU, Ahvaz, Iran. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0 license) (<http://creativecommons.org/licenses/by-nc/4.0/>).



گیاه پزشکی (مجله علمی کشاورزی)

جلد ۴۶، شماره ۳، پاییز ۱۴۰۲

doi 10.22055/ppr.2023.44689.1706

گزارش کوتاه انگلیسی

سموم حشره کش بالقوه مستخرج از برخی جدایه‌های قارچ‌های جنس *Akanthomyces*

تهمینه سلطانی^۱، فاطمه یاراحمدی^{۲*}، علی رجب پور^۲ و محمد حامد قدوم پاریزی پور^۳

- ۱- دانش‌آموخته کارشناسی ارشد حشره‌شناسی کشاورزی، گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه علوم کشاورزی و منابع طبیعی خوزستان، ملاتانی، ایران
- ۲- نویسنده مسوول: استاد، گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه علوم کشاورزی و منابع طبیعی خوزستان، ملاتانی، ایران (fa_yarahmadi@yahoo.com)
- ۳- استادیار، گروه گیاه‌پزشکی، دانشکده کشاورزی، دانشگاه علوم کشاورزی و منابع طبیعی خوزستان، ملاتانی، ایران

تاریخ پذیرش: ۱۴۰۲/۰۹/۰۴

تاریخ دریافت: ۱۴۰۲/۰۶/۱۰

چکیده

قارچ‌های بیمارگر حشرات عوامل کلیدی برای مدیریت آفات در کشاورزی پایدار می‌باشند. بسیاری از قارچ‌های بیمارگر حشرات می‌توانند متابولیت‌های ثانویه‌ای را تولید نمایند که دارای اثرات سمی هستند و در بیماری‌زایی آن‌ها نقشی اساسی دارند. در این مطالعه، این متابولیت‌ها در قارچ‌های بیمارگر حشرات شامل گونه *Akanthomyces lecanii* (جدایه‌های PAL6، PAL7 و PAL8) و گونه *A. muscarius* (جدایه AGM5) از نظر بیوشیمیایی با استفاده از کروماتوگرافی مایع با کارایی بالا (HPLC) مورد تجزیه و تحلیل قرار گرفتند. در میان زمان‌های مختلف بازدارنده مورد ثبت بین ۱-۱۳ دقیقه، چهار پیک مختلف در بین جدایه‌های PAL6، PAL7 و PAL8 مشاهده شد. در مورد جدایه GMA5، تنها یک پیک در زمان بازدارندگی ۱۱/۵۳ ثانیه دیده شد. تحلیل مقایسه‌ای نتایج نشان‌دهنده حضور پپتیدهای حلقوی سمی مانند Bassianolide در متابولیت‌های جدایه‌های PAL6، PAL7 و PAL8 در زمان بازدارندگی ۵ دقیقه و برای جدایه AGM5 در زمان بازدارندگی ۱۱/۵۳ دقیقه بود. این نتایج بیانگر تولید ترکیباتی سمی مانند Bassianolide توسط این قارچ‌های بیمارگر حشرات هستند که در ماهیت حشره‌کشی آن‌ها احتمالاً دخیل می‌باشند.

کلیدواژه‌ها: سموم قارچی، پپتیدهای حلقوی، حشره‌کش‌های زیستی، قارچ‌های بیمارگر حشرات

دبیر تخصصی: دکتر محسن یزدانیان

Citation: Soltani, T., Yarahmadi, F., Rajabpour, A. & Ghodoum Parizipour, M.H. (2023). Potential Insecticidal Toxins Extracted from Some Isolates of *Akanthomyces*. *Plant Protection (Scientific Journal of Agriculture)*, 46(3), 17-23. <https://doi.org/10.22055/ppr.2023.44689.1706>.